



Name: _____ Date: _____ Score: ____ / 8

CIRCLE THE CORRECT ANSWER FOR EACH QUESTION**Q1. What does DR ABC stand for in the primary survey?**

- A. Danger, Response, Airway, Breathing, Circulation
- B. Detect, React, Assess, Breathe, Compress
- C. Danger, Response, Assess, Breathe, Call
- D. Detect, Response, Airway, Bleeding, Circulation

Q2. How deep should chest compressions be for an adult?

- A. 2-3 cm
- B. 5-6 cm
- C. 8-10 cm
- D. As deep as possible

Q3. What is the correct compression-to-breath ratio for adult CPR?

- A. 15:2
- B. 30:2
- C. 5:1
- D. 10:2

Q4. At what rate should chest compressions be delivered?

- A. 60-80 per minute
- B. 80-100 per minute
- C. 100-120 per minute
- D. 120-140 per minute

Q5. What should you do if a casualty is unconscious but breathing normally?

- A. Begin CPR immediately
- B. Place in the recovery position
- C. Give 2 rescue breaths
- D. Call 999 and wait

Q6. For how long should you check for normal breathing?

- A. 5 seconds
- B. 10 seconds
- C. 20 seconds
- D. 30 seconds

Q7. What is the first action when treating a severe bleed?

- A. Apply a tourniquet
- B. Apply direct pressure to the wound
- C. Elevate the limb
- D. Call 999

Q8. What are the signs of a stroke? (FAST)

- A. Fever, Aching, Sweating, Tiredness
- B. Face drooping, Arm weakness, Speech difficulty, Time to call 999
- C. Faintness, Anxiety, Shaking, Trembling
- D. Flushing, Agitation, Swelling, Tightness

Answer Key (Trainer Use Only): Q1:A Q2:B Q3:B Q4:C
Q5:B Q6:B Q7:B Q8:B